

О. Лозинский, канд. хим. наук,
Т. Шокол, канд. хим. наук, shokol_tv@univ.kiev.ua,
Н. Горбуленко, канд. хим. наук,
В. Хилия, д-р хим. наук, чл.-корр. НАН Украины
КНУ имени Тараса Шевченко, Киев

НОВЫЙ МЕТОД СИНТЕЗА СИСТЕМЫ 4H,10H-ПИРАНО[2,3-f]ХРОМЕН-4,10-ДИОНА

Разработан новый метод синтеза системы 4H,10H-пирано[2,3-f]хромен-4,10-диона на основе циклизации 8-(3-диметиламино-2-пропеноил)-7-гидрокси-4H-хромен-4-она в уксусной кислоте.

Ключевые слова: 8-ацетил-7-гидроксихромоны, 7-гидрокси-8-(3-диметиламино-2-пропеноил)хромоны, диметилацеталь диметилформамида, 4H,10H-пирано[2,3-f]хромен-4,10-дионы.

O. Lozinski, PhD,
T. Shokol, PhD, shokol_tv@univ.kiev.ua,
N. Gorbulyenko, PhD,
V. Khilya, Dr. Sci., Corresponding Member of the NAS of Ukraine
Taras Shevchenko National University of Kyiv, Kyiv

A NOVEL METHOD OF SYNTHESIS OF THE 4H,10H-PYRANO[2,3-f]CHROMEN-4,10-DIONE SYSTEM

A simple and efficient method for the synthesis of 4H,10H-pyrano[2,3-f]chromen-4,10-dione system, which is characterized by rather high yields on relatively mild conditions was offered. Design of that system has been carried out by annelation of γ -pirone cycle to chromone core. 4H,10H-Pyrano[2,3-f]chromen-4,10-dione system was elaborated using 8-(3-dimethylamino-2-propenoyl)-7-hydroxy-4H-chromen-4-one cyclization in acetic acid. The starting enaminoketone derivative was synthesized in two stages from 7-acetoxychromone. 3-(4-Chlorophenyl)-6-ethyl-2-methyl-4-oxo-4H-7-chromenyl acetate was converted into 8-acetyl-3-(4-chlorophenyl)-6-ethyl-7-hydroxy-2-methyl-4H-4-chromenone by Fricke rearrangement. Subsequent heating of the latter with N,N-dimethylformamide dimethyl acetal in toluene gave the desired 3-(4-chlorophenyl)-8-(3-dimethylamino-2-propenoyl)-6-ethyl-7-hydroxy-4H-chromen-4-one. The advantage of such approach is the ability to obtain biopotent systems with different substituents in pyrone cycles, that will contribute to the search for new biologically active compounds. The structures of synthesized compounds have been assigned on the basis of analytical and spectra data.

Key words: 8-acetyl-7-hydroxychromones, 8-(3-dimethylamino-2-propenoyl)-7-hydroxychromones, N,N-dimethylformamide dimethyl acetal, 4H,10Hpyrano[2,3-f]chromen-4,10-diones.

УДК 547.814.5

T. Shokol, PhD, shokol_tv@univ.kiev.ua,
N. Gorbulyenko, PhD,
V. Khilya, Dr. Sci., Corresponding Member of the NAS of Ukraine
Taras Shevchenko National University of Kyiv, Kyiv

FEATURES OF THE α -AZAHETARYL-2-HYDROXYACETOPHENONES REACTION WITH CHLOROACETYL CHLORIDE

The reaction of α -azolyl-2-hydroxyacetophenones with chloroacetyl chloride in acetonitrile in the presence of pyridine resulted in 2-chloromethyl-3-azolylchromones, while both the α -(2-pyridyl) and α -(2-quinolyl) derivatives formed the products of the subsequent intramolecular cyclization with annelation of indolizine or pyrroloquinoline ring to the chromone core.

Key words: acylation, intramolecular cyclization, α -azahetaryl-2-hydroxyacetophenones, 3-azolyl-2-chloromethyl-7-hydroxy-4H-chromen-4-ones, 12H-chromeno[3,2-a]indolizin-12-ones, 7H-chromeno[3',2':3,4]pyrrolo[1,2-a]quinolin-7-ones.

Introduction. Isoflavones are naturally occurring products that possess a wide spectrum of biological activity [1]. 3-Hetarylchromones, heterocyclic analogs of natural isoflavones, are known for their inflammatory, antiviral, anabolic, analeptic, hypoglycemic and hypolipidemic activities [2]. α -Azahetaryl-2-hydroxyacetophenones are the key precursors for the synthesis of 3-azahetarylchromones. We have previously reported that α -azahetaryl-2-hydroxyacetophenones underwent acylation, followed by cyclization with acetic anhydride, trifluoroacetic anhydride and ethoxalyl chloride to give 2-substituted 3-azahetarylchromones [3]. Treatment of α -aryl-2-hydroxyacetophenones with chloroacetic anhydride gave 2-chloromethyl chromones [4].

As part of our ongoing interest in the synthesis of the new 3-azahetarylchromones and to extend our earlier work on acylation of α -azahetaryl-2-hydroxyacetophenones, the compounds **1.1a-1.10f** were treated with the excess of chloroacetyl chloride in acetonitrile in the presence of pyridine. Formation of 3-azolyl-7-chloroacetyl-2-chloromethylchromones **2.1a-2.6d** from α -azolyl-2-hydroxyacetophenones **1.1a-1.6d** occurs smoothly in 37-60 % yields. The ^1H NMR spectra of products **2.1a-2.6d** revealed the resonances of two methylene groups at 4.65-4.68 ppm (ClCH_2CO) and 5.10-5.58 ppm (2-ClCH_2) correspondingly.

Next we progressed on to elaboration of the hydrolysis of chloroacetates **2.1a-2.6d** to provide 3-azolyl-2-chloromethyl-7-hydroxychromones **3.1a-3.6d**. As expected, the ^1H NMR


spectra of compounds **3.1a-3.6d** showed disappearance of the signal at 4.65-4.68 ppm of the starting material, while 2-ClCH_2 signal remained intact. Finally, a new singlet at 10.77-10.96 ppm was assigned to the 7-OH group.

In contrast with α -azolyl derivatives, both the α -(2-pyridyl) (**1.7e-1.9e**) and α -(2-quinolyl) (**1.10f**) derivatives were found to form the products of the subsequent intramolecular cyclization with annelation of indolizine or pyrroloquinoline ring to the chromone core. This was confirmed by the presence of the characteristic signal of H6 at 7.70-7.80 ppm in the ^1H NMR spectra of the products **4.1e-4.3e** and H13 at 7.88 ppm in the spectrum of the product **6f**. Hydrolysis of compounds **4.1e, 4.2e** with 5 % NaOH afforded the 9-OH derivatives **5.1e, 5.2e**.

In conclusion we have reported that the resultant structures in reaction of α -azahetaryl-2-hydroxyacetophenone with chloroacetyl chloride relied upon the structure of heterocycle in starting material.

Experimental part. Reaction progress and identity of obtained compounds were monitored by TLC on Merc 60 F₂₅₄ silica gel plates using $\text{CHCl}_3\text{-MeOH}$ (9:1) system. NMR spectra were recorded on Mercury-400 spectrometer (spectrometer frequency for ^1H : 400 MHz) from $\text{DMSO-}d_6$ and CDCl_3 solns. The TMS signal was used as an internal standart. Elemental analyses for C, H, and N were performed using Perkin-Elmer C, H, N Analyser.

Compounds **2.1a, 2.2a, 2.4b, 2.5c, 3.1a, 3.2a** and **3.4b** were synthesized according to a procedure reported in the literature [5-7].



1.1a, 1.5c R=H, Alk=5-Et; **1.2a, 1.3b** R=H, Alk=5-Pr; **1.4b** R=H, Alk=5-Hex; **1.6d** R=H, Alk=3-Me; **1.7e** R=Ac, Alk=3-Me; **1.8e** R=Ac, Alk=5-Et; **1.9e** R=H, Alk=5-Pr; **1.10f** R=Me, Alk=5-Et; **2.1a, 3.1a, 2.5c, 3.5c** Alk=6-Et; **2.2a, 3.2a, 2.3b, 3.3b** Alk=6-Pr; **2.4b, 3.4b** Alk=6-Hex; **2.6d, 3.6d** Alk=8-Me; **4.1e** R=Ac, Alk=8-Me; **4.2e** R=Ac, Alk=10-Et; **4.3e** R=ClCH₂CO, Alk=10-Pr; **5.1e** R=H, Alk=8-Me; **5.2e** R=H, Alk=10-Et

3-Azoly-2-chloromethyl-4-oxo-4H-7-chromenyl 2-chloroacetates (2.3b, 2.6d). General procedure. Pyridine (1.22 mL, 15 mmol) and chloroacetyl chloride (1.20 mL, 15 mmol) were added to a solution of compound **1.3b** or **1.6d** (5 mmol) in acetonitrile (15 mL), held for 24 h at room temperature, and obtained precipitate was filtered off and washed with water.

3-(1,3-Benzothiazol-2-yl)-2-chloromethyl-4-oxo-6-propyl-4H-7-chromenyl 2-chloroacetate (2.3b). Yellow solid; yield: 1.39 g (60 %); mp 158 °C. ¹H NMR (DMSO-d₆): δ=0.98 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 1.66 (m, 2H, CH₂), 2.68 (t, ³J_{HH}=7.2 Hz, 2H, CH₂), 4.65 (s, 2H, ClCH₂CO), 5.58 (s, 2H, CH₂Cl), 7.46 (t, ³J_{HH}=7.2 Hz, 1H, H6'), 7.53 (t, ³J_{HH}=7.2 Hz, 1H, H5'), 7.67 (s, 1H, H8), 8.07 (d, ³J_{HH}=8.4 Hz, 1H, H7'), 8.09 (d, ³J_{HH}=8.4 Hz, 1H, H4'), 8.13 (s, 1H, H5). Anal. Calcd for C₂₂H₁₇Cl₂NO₄S: C, 57.15; H, 3.71; Cl, 15.34; N, 3.03; S, 6.93. Found: C, 57.41; H, 3.87; Cl, 15.46; N, 2.76; S, 7.11.

2-Chloromethyl-8-methyl-3-(2-methyl-1,3-thiazol-4-yl)-4-oxo-4H-7-chromenyl 2-chloroacetate (2.6d). Colorless solid; yield: 0.73 g (37%); mp 195 °C. ¹H NMR (DMSO-d₆): δ=2.37 (s, 3H, 8-CH₃), 2.76 (s, 3H, 2'-CH₃), 4.68 (s, 2H, ClCH₂CO), 5.10 (s, 2H, CH₂Cl), 7.29 (d, ³J_{HH}=8.4 Hz, 1H, H6'), 7.94 (s, 1H, H5'), 8.02 (d, ³J_{HH}=8.4 Hz, 1H, H5). Anal. Calcd for C₁₇H₁₃Cl₂NO₄S: C, 51.27; H, 3.29; Cl, 17.80; N, 3.52; S, 8.05. Found: C, 51.49; H, 3.42; Cl, 17.92; N, 3.20; S, 8.23.

3-Azoly-2-chloromethyl-7-hydroxy-4H-chromen-4-ones (3.3b, 3.5c, 3.6d). General procedure. A solution of compound **2.3b**, **2.5c** or **2.6d**, (3 mmol) in a mixture of EtOH (50 mL) and 37% HCl (1 mL) was refluxed for 1.5 h, then set aside overnight at room temperature, filtered off and washed with ethanol and water.

3-(1,3-Benzothiazol-2-yl)-2-chloromethyl-7-hydroxy-6-propyl-4H-4-chromenone (3.3b). Yellow solid; yield:

1.11 g (96 %); mp 305 °C. ¹H NMR (DMSO-d₆): δ=1.00 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 1.67 (m, 2H, CH₂), 2.65 (t, ³J_{HH}=7.2 Hz, 2H, CH₂), 5.56 (s, 2H, CH₂Cl), 6.95 (s, 1H, H8), 7.43 (t, ³J_{HH}=7.6 Hz, 1H, H6'), 7.51 (t, ³J_{HH}=7.6 Hz, 1H, H5'), 7.86 (s, 1H, H5), 8.04 (d, ³J_{HH}=8.0 Hz, 1H, H7'), 8.07 (d, ³J_{HH}=8.0 Hz, 1H, H4'), 10.89 (s, 1H, OH). Anal. Calcd for C₂₀H₁₆ClNO₃S: C, 62.26; H, 4.18; Cl, 9.19; N, 3.63; S, 8.31. Found: C, 62.48; H, 4.43; Cl, 8.99; N, 3.33; S, 8.54.

2-Chloromethyl-6-ethyl-7-hydroxy-3-(5-phenyl-1,3,4-thiadiazol-2-yl)-4H-4-chromenone (3.5c). Colorless solid; yield: 1.19 g (99 %); mp 236 °C. ¹H NMR (DMSO-d₆): δ=1.28 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 2.71 (q, ³J_{HH}=7.2 Hz, 2H, CH₂), 5.49 (s, 2H, CH₂Cl), 6.98 (s, 1H, H8), 7.55 (m, 3H, H3', H4', H5'), 7.89 (s, 1H, H5), 8.07 (m, 2H, H2', H6'), 10.96 (s, 1H, OH). Anal. Calcd for C₂₀H₁₅ClN₂O₃S: C, 60.23; H, 3.79; Cl, 8.89; N, 7.02; S, 8.04. Found: C, 60.45; H, 3.86; Cl, 9.00; N, 6.89; S, 7.79.

2-Chloromethyl-7-hydroxy-8-methyl-3-(2-methyl-1,3-thiazol-4-yl)-4H-4-chromenone (3.6d). Colorless solid; yield: 0.95 g (98 %); mp 240 °C. ¹H NMR (DMSO-d₆): δ=2.30 (s, 3H, 8-CH₃), 2.79 (s, 3H, 2'-CH₃), 5.02 (s, 2H, CH₂Cl), 7.02 (d, ³J_{HH}=8.8 Hz, 1H, H6'), 7.78 (d, ³J_{HH}=8.8 Hz, 1H, H5'), 7.91 (s, 1H, H5'), 10.77 (s, 1H, OH). Anal. Calcd for C₁₅H₁₂ClNO₃S: C, 55.99; H, 3.76; Cl, 11.02; N, 4.35; S, 9.96. Found: C, 55.73; H, 3.85; Cl, 11.00; N, 4.60; S, 9.79.

12-Oxo-12H-chromeno[3,2-a]indolizin-9-yl acetates (4.1e, 4.2e). General procedure. Pyridine (1.6 mL, 20 mmol) and chloroacetyl chloride (0.88 g, 11 mmol) were added to a solution of compound **1.7e** or **1.8e** (10 mmol) in acetonitrile (15 mL). The reaction mixture was refluxed for 20 min and then was cooled to room temperature. The resultant precipitate was filtered off and recrystallized from *o*-xylene.

8-Methyl-12-oxo-12H-chromeno[3,2-a]indolizin-9-yl acetate (4.1e). Green solid; yield: 2.46 g (80 %); mp 265 °C. ¹H NMR (DMSO-d₆): δ=2.33 (s, 3H, CH₃), 2.37

(s, 3H, CH₃CO), 7.05 (d, ³J_{HH}=7.6 Hz, 1H, H10), 7.07 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.32 (t, ³J_{HH}=7.6 Hz, 1H, H2), 7.74 (s, 1H, H6), 8.07 (d, ³J_{HH}=8.4 Hz, 1H, H11), 8.25 (d, ³J_{HH}=8.4 Hz, 1H, H1), 8.61 (d, ³J_{HH}=6.4 Hz, 1H, H4). Anal. Calcd for C₁₈H₁₃NO₄: N, 4.56. Found: N, 4.45.

10-Ethyl-12-oxo-12H-chromeno[3,2-a]indolizin-9-yl acetate (4.2e). Green solid; yield: 1.99 g (62 %); mp 202 °C. ¹H NMR (DMSO-d₆): δ=1.25 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 2.37 (s, 3H, CH₃CO), 2.65 (q, ³J_{HH}=7.2 Hz, 2H, CH₂), 7.08 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.33 (t, ³J_{HH}=7.6 Hz, 1H, H2), 7.40 (s, 1H, H8), 7.70 (s, 1H, H6), 8.08 (s, 1H, H11), 8.25 (d, ³J_{HH}=8.4 Hz, 1H, H1), 8.60 (d, ³J_{HH}=6.4 Hz, 1H, H4). Anal. Calcd for C₁₉H₁₅NO₄: N, 4.36. Found: N, 4.21.

12-Oxo-10-propyl-12H-chromeno[3,2-a]indolizin-9-yl 2-chloroacetate (4.3e) was obtained from comp. **1.9e** (2.71 g, 10 mmol) and chloroacetyl chloride (1.75 mL, 22 mmol) according to a procedure for comp. **4.1e**. Green solid; yield: 0.25 g (68 %); mp 278 °C. ¹H NMR (DMSO-d₆): δ=0.92 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 1.66 (m, 2H, CH₂), 2.62 (t, ³J_{HH}=7.2 Hz, 2H, CH₂), 4.62 (s, 2H, ClCH₂CO), 7.07 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.33 (t, ³J_{HH}=7.6 Hz, 1H, H2), 7.40 (s, 1H, H8), 7.74 (s, 1H, H6), 8.08 (s, 1H, H11), 8.28 (d, ³J_{HH}=8.0 Hz, 1H, H1), 8.62 (d, ³J_{HH}=5.6 Hz, 1H, H4). Anal. Calcd for C₂₀H₁₆ClNO₄: Cl, 9.59; N, 3.79. Found: Cl, 9.59; N, 3.84.

9-Hydroxy-12H-chromeno[3,2-a]indolizin-12-ones (5.1e, 5.2e). General procedure. A mixture of the corresponding acetates **4.1e** or **4.2e** (5 mmol) in 50 mL of EtOH and 5% water solution of NaOH (4 mL, 5 mmol) was refluxed for 1 h, diluted with water (100 mL), refluxed for 5-10 min, neutralized with HCl to pH 7 and the resultant precipitate was filtered off and recrystallized from DMF.

9-Hydroxy-8-methyl-12H-chromeno[3,2-a]indolizin-12-one (5.1e). Yellow solid; yield: 1.13 g (86 %); mp 218 °C. ¹H NMR (DMSO-d₆): δ=2.30 (s, 3H, CH₃), 7.10 (d, ³J_{HH}=8.4 Hz, 1H, H10), 7.30 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.70 (s, 1H, H6), 7.73 (t, ³J_{HH}=7.6 Hz, 1H, H2), 8.08 (d, ³J_{HH}=8.4 Hz, 1H, H11), 8.25 (d, ³J_{HH}=7.6 Hz, 1H, H1), 8.60 (d, ³J_{HH}=6.4 Hz, 1H, H4), 10.40 (s, 1H, OH). Anal. Calcd for C₁₆H₁₁NO₃: N, 5.28. Found: N, 5.09.

10-Ethyl-9-hydroxy-12H-chromeno[3,2-a]indolizin-12-one (5.2e). Yellow solid; yield: 1.00 g (72 %); mp 217 °C

¹H NMR (DMSO-d₆): δ=1.25 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 2.65 (q, ³J_{HH}=7.2 Hz, 2H, CH₂), 6.88 (s, 1H, H8), 7.35 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.73 (t, ³J_{HH}=7.6 Hz, 1H, H2), 7.80 (s, 1H, H6), 8.09 (s, 1H, H11), 8.25 (d, ³J_{HH}=8.4 Hz, 1H, H1), 8.59 (d, ³J_{HH}=6.4 Hz, 1H, H4), 10.41 (s, 1H, OH). Anal. Calcd for C₁₇H₁₃NO₃: N, 5.02. Found: N, 5.00.

9-Ethyl-10-methoxy-7H-chromeno[3',2':3,4]-pyrrolo[1,2-a]quinolin-7-one (6f). Method A. Product **6f** was obtained from compound **1.10f** (0.48 g, 1.5 mmol), pyridine (0.3 mL, 3.7 mmol) and chloroacetyl chloride (0.13 mL, 1.65 mmol) in CH₃CN (11 mL) according to a procedure for compound **4.1e**.

Method B. Product **6f** was obtained from compound **1.10f** (0.64 g, 2 mmol), pyridine (0.3 mL, 3.7 mmol) and chloroacetic anhydride (0.88 g, 10 mmol) in CH₃CN (10 mL). The reaction mixture was refluxed for 5-10 min, poured onto water (100 mL) and the resultant precipitate was filtered off and washed with EtOH. Yellow solid; yield: 0.27 g (52 %, method A), 0.5 g (74 %, method B); mp 301-302 °C. ¹H NMR (CDCl₃): δ=1.26 (t, ³J_{HH}=7.2 Hz, 3H, CH₃), 2.72 (q, ³J_{HH}=7.2 Hz, 2H, CH₂), 3.94 (s, 3H, CH₃O), 6.84 (s, 1H, H11), 7.47 (t, ³J_{HH}=7.6 Hz, 1H, H2), 7.51 (d, ³J_{HH}=8.4 Hz, 1H, H4), 7.65 (t, ³J_{HH}=7.6 Hz, 1H, H3), 7.82 (d, ³J_{HH}=7.6 Hz, 1H, H1), 7.88 (s, 1H, H13), 8.13 (s, 1H, H8), 7.96 (d, ³J_{HH}=8.8 Hz, 1H, H5), 8.37 (d, ³J_{HH}=8.8 Hz, 1H, H6). Anal. Calcd for C₂₂H₁₇NO₃: N, 4.08. Found: N, 4.01.

References

1. Казаков А.Л., Хиля В.П., Мережицкий В.В., Литкей Ю. Природные и модифицированные изофлавоноиды. Ростов-на-Дону, Издательство Ростовского университета, 1985, 184 с.
2. Kazakov A.L., Khilya V.P., Mezheritskii V.V., Litkei G. Natural and modified isoflavonoids. Rostov-on-the-Don, Rostovsk. Univ., 1985, 184 p. (in Russian).
3. Горбуленко Н.В., Хиля В.П. Укр. хим. журнал, 1994, 60(1), 79-91.
4. Gorbulyenko N.V., Khilya V.P. Ukrainskii khimicheskii zhurnal, 1994, 60(1), 79-91.
5. Frasinuk M.S., Khilya V.P. Chem. Heterocycl. Comp., 1999, 35(1), 3-22.
6. Cassella Farbwerke Mainkur A.-G. GB Pat. Appl. 1016088, 1966.
7. Shokol T.V., Gorbulyenko N.V., Khilya V.P. Chem. Heterocycl. Comp., 2012, 47(10), 1298-1299.
8. Shokol T.V., Gorbulyenko N.V., Turov A.V., Khilya V.P. Chem. Heterocycl. Comp., 2012, 48(8), 1181-1186.
9. Shokol T.V., Semenyuchenko V.V., Khilya V.P. Chem. Heterocycl. Comp., 2004, 40(12), 1588-1594.

Надійшла до редколегії 30.10.14

Т. Шокіл, канд. хім. наук, shokol_tv@univ.kiev.ua,
Н. Горбуленко, канд. хім. наук,
В. Хиля, д-р хім. наук, чл.-кор. НАН України
КНУ імені Тараса Шевченка, Київ

ОСОБЛИВОСТІ РЕАКЦІЇ α-АЗАГЕТАРИЛ-2-ГІДРОКСИАЦЕТОФЕНОНІВ З ХЛОРАЦЕТИЛХЛОРИДОМ

Встановлено, що реакція α-азоліл-2-гідроксиацетофенонів з хлорацетилхлоридом в ацетонітрилі в присутності піридину приводить до 2-хлорметил-3-азолілхромонів, тоді як α-(2-піридил)- та α-(2-хіноліл)- похідні утворюють продукти внутрішньомолекулярної циклізації з анелюванням індолізинового або піролохінолінового циклу до хромонового ядра.

Ключові слова: ацилювання, внутрішньомолекулярна циклізація, α-азагетарил-2-гідроксиацетофенони, 3-азоліл-2-хлорметил-7-гідрокси-4Н-хромен-4-они, 12Н-хромено[3,2-а]індолізин-12-они, 7Н-хромено[3',2':3,4]піроло[1,2-а]хінолін-7-они.

Т. Шокіл, канд. хім. наук, shokol_tv@univ.kiev.ua,
Н. Горбуленко, канд. хім. наук,
В. Хиля, д-р хім. наук, чл.-корр. НАН України
КНУ імені Тараса Шевченка, Київ

ОСОБЕННОСТИ РЕАКЦИИ α-АЗАГЕТАРИЛ-2-ГИДРОКСИАЦЕТОФЕНОНОВ С ХЛОРАЦЕТИЛХЛОРИДОМ

Установлено, что реакция α-азоліл-2-гідроксиацетофенонов с хлорацетилхлоридом в ацетонитриле в присутствии пиридина приводит к 2-хлорметил-3-азолілхромонам, тогда как α-(2-пиридил)- и α-(2-хиноліл)- производные образуют продукты внутримолекулярной циклизации с анелюванием индолизинового или пиролохинолинового цикла к хромоновому ядру.

Ключевые слова: ацилирование, внутримолекулярная циклизация, α-азагетарил-2-гідроксиацетофеноны, 3-азоліл-2-хлорметил-7-гідрокси-4Н-хромен-4-оны, 12Н-хромено[3,2-а]індолізин-12-оны, 7Н-хромено[3',2':3,4]піроло[1,2-а]хінолін-7-оны.