

UDC 546.3-386:547.771.057

DOI: [https://doi.org/10.17721/1728-2209.2025.1\(60\).10](https://doi.org/10.17721/1728-2209.2025.1(60).10)

Yuliia DAVYDENKO, PhD (Chem.)

ORCID ID: 0000-0003-0223-0363

e-mail: [y.davydenko@knu.ua](mailto:y.davydenko@knu.ua)

Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

## SYNTHESIS AND CHARACTERIZATION OF COORDINATION COMPOUNDS OF TRANSITION METALS BASED ON 5-METHYL-3-(TRIFLUOROMETHYL)-1H-PYRAZOLE

**Background.** The design of coordination compounds using pyrazole-based ligands is of fundamental importance in contemporary materials chemistry due to their versatile coordination modes and diverse applications. Introducing trifluoromethyl groups onto such ligands is a powerful strategy to modulate the electronic and steric properties of the resulting metal complexes, significantly influencing their acidity, stability, and reactivity. In this context, 5-methyl-3-(trifluoromethyl)-1H-pyrazole stands out as a particularly promising ligand whose coordination chemistry with transition metals remains surprisingly underexplored. Despite its potential, a systematic investigation into the synthesis, structural diversity, and properties of its coordination compounds is largely absent from recent literature. This gap presents a missed opportunity, as the unique electronic profile of this ligand could unlock new functionalities in molecular materials. To address this, we present a comprehensive study on the synthesis and characterization of a new family of coordination compounds of first-row transition metals (Mn, Co, Ni, Cu, Zn) with the 5-methyl-3-(trifluoromethyl)-1H-pyrazole ligand. This work aims to systematically explore its coordination landscape, providing fundamental insights for the future design of advanced functional materials.

**Methods.** In this work, the coordination compounds with transition metals were obtained using the 5-methyl-3-(trifluoromethyl)-1H-pyrazole ligand. A variety of techniques were used to identify and characterize the complexes and the ligand, including infrared, UV/Vis, and NMR spectroscopy, as well as microanalyses.

**Results.** The interaction of 3d metals such as Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) with 5-methyl-3-(trifluoromethyl)-1H-pyrazole in non-aqueous solutions was found to be solvent-dependent. In a solvent environment CH<sub>3</sub>CN and CH<sub>3</sub>OH (M(Ac)<sub>2</sub>·4H<sub>2</sub>O–L–CH<sub>3</sub>OH/CH<sub>3</sub>CN systems) at M:L ratios of 1:1 and 1:2, products with the composition M(Ac)<sub>2</sub>·2L were formed. Meanwhile, in dimethylformamide (M(Ac)<sub>2</sub>·4H<sub>2</sub>O–L–DMF systems) under the same ratios, the complex Mn(Ac)<sub>2</sub>·L·DMF, which incorporated a solvent molecule, was isolated.

**Conclusions.** An organic ligand, 5-methyl-3-(trifluoromethyl)-1H-pyrazole, was synthesized. The interaction of 3d metals (M = Mn, Co, Ni, Cu, Zn) M(Ac)<sub>2</sub>·xH<sub>2</sub>O with 5-methyl-3-(trifluoromethyl)-1H-pyrazole in non-aqueous (CH<sub>3</sub>CN, CH<sub>3</sub>OH and DMF) solutions was investigated. Methods for the synthesis of Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) compounds with 5-methyl-3-(trifluoromethyl)-1H-pyrazole were developed. Based on the data of elemental analysis and methods of IR, NMR, and electronic spectroscopy, the composition and structure of the obtained complexes have been proposed. It has been shown that in the obtained complexes: the most typical method of coordination of the pyrazole cycle is realized: monodentate - through the pyridine nitrogen atom; the formation of two types of mononuclear Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) complexes is observed: M(Ac)<sub>2</sub>·2L and M(Ac)<sub>2</sub>·L·DMF; in the obtained complexes M(Ac)<sub>2</sub>·2L and M(Ac)<sub>2</sub>·L·DMF complexes, a six-coordinate environment of the central atom is realized due to four oxygen atoms from bidentate chelate coordinated acetate groups and two nitrogen atoms from molecules of a non-deprotonated ligand.

**Keywords:** 5-methyl-3-(trifluoromethyl)-1H-pyrazole, complexes, 3d metals, <sup>1</sup>H NMR spectroscopy, IR spectroscopy.

### Background

The design and synthesis of coordination compounds using nitrogen-containing heterocyclic ligands are of fundamental importance in contemporary inorganic and materials chemistry. Among these, pyrazole-based ligands have been extensively employed to construct a diverse range of metal complexes with fascinating structural topologies and functional properties (Davydenko et al., 2024). The versatility of the pyrazole core, with its two adjacent nitrogen atoms, allows for various coordination modes-acting as a monodentate ligand (Davydenko et al., 2023; Vynohradov et al., 2022), a bridging unit to form polynuclear clusters (Davydenko, 2012; 2022; 2024; 2025;), or as a component of more complex chelating systems (Mezei et al., 2007; Pandolfo et al., 2017). This adaptability has led to the development of coordination compounds with remarkable applications in catalysis (Annes et al., 2022; Titi et al., 2023), magnetism (Li et al., 2023; Wang et al., 2023), luminescent sensing (Sun et al., 2023; Zhang et al., 2024), and the construction of metal-organic frameworks (MOFs) (Roy et al., 2023).

The strategic introduction of fluorine atoms or trifluoromethyl groups onto organic ligands has emerged as a powerful tool to modulate the electronic and steric properties of the resulting metal complexes. The strong electron-withdrawing nature of the CF<sub>3</sub> group can significantly influence the acidity of the N-H proton of the pyrazole ring, the donor ability of the nitrogen atoms, and the overall stability and reactivity of the coordination

compounds. Furthermore, the presence of fluorine can enhance the volatility, thermal stability, and solubility of the complexes in non-polar solvents, and can lead to unique solid-state packing through non-covalent interactions. In this context, 5-methyl-3-(trifluoromethyl)-1H-pyrazole stands out as a particularly promising ligand. The CF<sub>3</sub> group is expected to significantly lower the pK<sub>a</sub> of the pyrazole proton, thereby facilitating deprotonation and the formation of pyrazolate-based complexes. Moreover, the electronic and steric effects of CF<sub>3</sub> substituents are anticipated to impart unique characteristics to the resulting coordination compounds, potentially leading to novel properties.

Despite the considerable potential of 5-methyl-3-(trifluoromethyl)-1H-pyrazole as a ligand, a survey of recent literature reveals that the coordination chemistry of this ligand with a broad range of transition metals remains surprisingly underexplored. Foundational studies have demonstrated the ability of fluorinated pyrazoles to form stable complexes (Fustero et al., 2014; Mykhailiuk, 2020), and recent research has touched upon related systems for applications such as photoswitching (Park et al., 2020; Kieffer et al., 2022; Kolarski et al., 2020) and catalysis with other fluorinated ligands (Sánchez-Cantú et al., 2018; Titi et al., 2023). However, a systematic investigation into the synthesis, structural diversity, and functional properties of coordination compounds based on 5-methyl-3-(trifluoromethyl)-1H-pyrazole in the last 5 years is largely absent. This represents a significant gap in the field, as the unique electronic profile of this ligand could unlock new

© Davydenko Yuliia, 2025

opportunities in the design of functional molecular materials. For instance, the electron-poor nature of the pyrazolate ring could influence the magnetic exchange interactions in polynuclear complexes or tune the redox potentials of catalytically active metal centers.

Herein, we present a comprehensive study on the synthesis and characterization of a new family of coordination compounds of first-row transition metals (Mn, Co, Ni, Cu, Zn) with the 5-methyl-3-(trifluoromethyl)-1H-pyrazole ligand.

This work aims to systematically explore the coordination landscape of 5-methyl-3-(trifluoromethyl)-1H-pyrazole, providing fundamental insights that will pave the way for the future design of advanced functional materials based on this versatile building block.


### Methods

All chemicals and solvents were commercial products of reagent grade and used without further purification. Microanalyses were performed with a Perkin-Elmer 2400 CHN. IR spectra (KBr pellets) were recorded with a Perkin-

Elmer Spectrum BX FT-IR in the range of 400–4000  $\text{cm}^{-1}$ . Absorbance UV/Vis spectra were registered with a Varian Cary 50 spectrometer in the range of 200–1000 nm at room temperature.  $^1\text{H}$  NMR spectra were recorded on a Bruker AC-400 spectrometer (400 MHz) at room temperature.

### Results

*The synthesis of 5-methyl-3-(trifluoromethyl)-1H-pyrazole (L).* The synthesis of 5-methyl-3-(trifluoromethyl)-1H-pyrazole was carried out according to the following scheme 1. 12.95 g (0.084 mol) of 1,1,1-trifluoro-2,4-pentadiol in 110 mL of ethanol was added dropwise, with constant stirring, 4.1 mL (4.2 g; 0.084 mol) of hydrazine hydrate. The reaction mixture was stirred for 3 hours while cooling on an ice bath. After complete addition of the hydrazine hydrate solution, the reaction mixture was left overnight. The solvents were completely distilled off on a rotary evaporator. The product was obtained in the form of light yellow crystals, which are highly soluble in DMF, acetonitrile, and methanol.



Scheme 1. Synthesis of 5-methyl-3-(trifluoromethyl)-1H-pyrazole

The yield of the product was 77 %.  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 2.30 (s., 3H;  $\text{CH}_3$ ), 6.22 (s., 1H; pz-CH), 13.07 (br. s., 1H; NH). Elemental analysis: calculated (%): C 40.01; H 3.36; N 18.66. Found: C 40.04; H 3.35; N 18.70. IR (KBr disk,  $\text{cm}^{-1}$ , selected bands): 3194s, 3120s, 2890s, 2814m, 2562w, 1587s, 1504s, 1254s, 1159s.

*The general method of the synthesis of the coordination compounds.* The synthesis of coordination compounds with 5-methyl-3-(trifluoromethyl)-1H-pyrazole (L) was performed with  $\text{M}(\text{CH}_3\text{COO})_2 \cdot x\text{H}_2\text{O}$  of different 3d-metals (scheme 2).



L – 5-methyl-3-(trifluoromethyl)-1H-pyrazole.

M = Mn, Zn, Cu, Co, Ni.

Solv –  $\text{CH}_3\text{CN}$ , DMF,  $\text{CH}_3\text{OH}$ .

Scheme 2. Synthesis of the coordination compounds

The synthesis of coordination compounds was carried out in a flat-bottomed reactor with a volume of 25 mL. A pre-dissolved salt (0.001 mol; 0.002 mol) was added to the reactor, to which a pre-dissolved ligand (0.001 mol; 0.002 mol) was slowly added dropwise. Metal : ligand ratios of both 1:1 and 1:2 were used. The amount of solvent was varied between 2 and 9 ml, depending on the nature of the solvent and the solubility of the reagents. The reaction mixture was stirred on a magnetic stirrer for 15 minutes at a temperature of 21–60  $^\circ\text{C}$  until the complete conversion of the starting reagents. When an insoluble reaction product was formed, it was filtered off and dried in air. If a product in the form of a solution was formed, it was left to crystallize slowly in air at room temperature. The composition of the products obtained from the interaction of 5-methyl-3-(trifluoromethyl)-1H-pyrazole with non-aqueous solutions of 3d-metal acetates (Mn, Zn, Cu, Co, and Ni) and the elemental analysis data are presented in tabl. 1.

### Discussion and conclusions

Thus, as a result of the interaction between 5-methyl-3-(trifluoromethyl)-1H-pyrazole and non-aqueous solutions of 3d-metal acetates (Mn, Zn, Cu, Co, Ni), different products were obtained. Specifically, in  $\text{M}(\text{Ac})_2 \cdot 4\text{H}_2\text{O} - \text{L} - \text{CH}_3\text{OH}/\text{CH}_3\text{CN}$



systems with M:L ratios of 1:1 and 1:2, compounds with the composition  $\text{M}(\text{Ac})_2 \cdot 2\text{L}$  were formed (fig. 1, a). In contrast, in the systems with dimethylformamide ( $\text{M}(\text{Ac})_2 \cdot 4\text{H}_2\text{O} - \text{L} - \text{DMF}$ ) at similar ratios,  $\text{Mn}(\text{Ac})_2 \cdot \text{L} - \text{DMF}$  was obtained (fig. 1, b), where the solvent became part of the complex. These conclusions are based on data from elemental analysis, IR, UV/Vis, and NMR spectroscopy.

*NMR spectroscopy.*  $^1\text{H}$  NMR spectra of the ligand ( $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 2.30 (s., 3H;  $\text{CH}_3$ ), 6.22 (s., 1H; pz-CH), 13.07 (br. s., 1H; NH)) (fig. 2), as a reference sample, and compound obtained as a result of synthesis  $\text{Zn}(\text{Ac})_2 \cdot 2\text{H}_2\text{O} - \text{L} - \text{Solv}$  (fig. 3), demonstrate the fact of complex formation.  $^1\text{H}$  NMR spectrum of the complex ( $\text{DMSO-d}_6$ ) indicates the presence of the peaks are observed at  $\delta$ : 2.30 and 3.10 (two singlets corresponding to three hydrogen atoms from two metal groups, two singlets at  $\delta$ : 6.24 and 6.26 from two 1H pyrazole nuclei and a broadened singlet at 13.07 from the 1H group NH (fig. 3), which indicates the formation of a mixture of two zinc complexes. This is associated with a change in the position of the  $\text{CH}_3$  and  $\text{CF}_3$  substituents in the 3rd and 5th positions of the pyrazole nucleus.

*IR spectroscopy.* In the comparative analysis of the IR spectra of the synthesized complexes and ligand, the main attention was paid to the absorption bands corresponding to the oscillations of the functional groups of the ligand (IR (KBr disk,  $\text{cm}^{-1}$ , selected bands): 3194s, 3120s, 2890s, 2814m, 2562w, 1587s, 1504s, 1254s, 1159s). In the IR spectra of synthesized compounds of general composition  $\text{M}(\text{Ac})_2 \cdot 2\text{L}$  and  $\text{M}(\text{Ac})_2 \cdot \text{L} - \text{DMF}$ , absorption bands are observed in the region of 3300–3200  $\text{cm}^{-1}$ , which is characteristic of valence vibrations of the NH group and indicates the presence of unprotonated molecules of the ligand 5-methyl-3-(trifluoromethyl)-1H-pyrazole. The following bands indicate the presence of acetate groups:  $\nu_{\text{as}}(\text{C}=\text{O}) = 1510\text{--}1520 \text{ cm}^{-1}$ ,  $\nu_{\text{s}}(\text{C}-\text{O}) = 1450\text{--}1460 \text{ cm}^{-1}$  (fig. 4, 5). In  $\text{M}(\text{Ac})_2 \cdot \text{L} - \text{DMF}$  complexes, the presence of a narrow intense absorption band at 1640  $\text{cm}^{-1}$  attributed to the valence vibrations of the CO group indicates the coordination of dimethylformamide molecules (fig. 6).

Table 1

The original system	M:L:Solv	Composition of the extracted product	C, % calc./exp.	H, % calc./exp.	N, % calc./exp.
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,78/34,52	2,89/2,86	10,14/10,08
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> OH	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,78/35,12	2,89/2,83	10,14/10,04
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- DMF	1 : 1: 9mL	M(Ac) <sub>2</sub> ×L×DMF	35,82/35,23	3,88/3,83	12,53/12,31
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,78/34,17	2,89/2,84	10,14/9,87
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> OH	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,78/34,56	2,89/2,87	10,14/10,02
Mn(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- DMF	1 : 2: 11mL	M(Ac) <sub>2</sub> ×L×DMF	35,82/34,89	3,88/3,81	12,53/12,20
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O- L- CH <sub>3</sub> OH	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	33,56/33,13	2,79/2,64	9,79/9,55
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	33,56/33,28	2,79/2,58	9,79/9,12
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O-L- DMF	1 : 1: 9mL	M(Ac) <sub>2</sub> ×L×DMF	34,78/34,34	3,76/3,59	12,17/11,49
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	33,56/33,12	2,79/2,66	9,79/9,47
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O- L- CH <sub>3</sub> OH	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	33,56/33,43	2,79/2,54	9,79/9,52
Zn(Ac) <sub>2</sub> ×2H <sub>2</sub> O-L- DMF	1 : 2: 11mL	M(Ac) <sub>2</sub> ×L×DMF	34,78/33,28	3,76/3,22	12,17/12,01
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-CH <sub>3</sub> CN	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	33,74/32,98	2,81/2,56	9,84/9,51
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-CH <sub>3</sub> OH	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	33,74/33,15	2,81/2,67	9,84/9,26
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-DMF	1 : 1: 9mL	M(Ac) <sub>2</sub> ×L×DMF	34,93/34,58	3,78/3,41	12,22/12,00
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-CH <sub>3</sub> CN	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	33,74/33,46	2,81/2,48	9,84/9,56
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-CH <sub>3</sub> OH	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	33,74/33,27	2,81/2,56	9,84/9,49
Cu(Ac) <sub>2</sub> ×H <sub>2</sub> O-L-DMF	1 : 2: 11mL	M(Ac) <sub>2</sub> ×L×DMF	34,93/34,26	3,78/3,63	12,22/11,98
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L-CH <sub>3</sub> CN	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,28/33,98	2,85/2,78	10,00/9,68
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L-CH <sub>3</sub> OH	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,28/33,82	2,85/2,81	10,00/9,74
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L-DMF	1 : 1: 9mL	M(Ac) <sub>2</sub> ×L×DMF	35,39/34,91	3,83/3,76	12,38/12,23
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,28/34,02	2,85/2,74	10,00/9,84
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> OH	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,28/33,83	2,85/2,68	10,00/9,76
Co(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- DMF	1 : 2: 11mL	M(Ac) <sub>2</sub> ×L×DMF	35,39/33,78	3,83/3,71	12,38/12,23
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,28/33,96	2,85/2,78	10,00/9,79
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> OH	1 : 1: 9mL	M(Ac) <sub>2</sub> ×2L	34,28/33,81	2,85/2,81	10,00/9,68
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- DMF	1 : 1: 9mL	M(Ac) <sub>2</sub> ×L×DMF	35,39/34,99	3,83/3,69	12,38/12,02
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> CN	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,28/33,46	2,85/2,71	10,00/9,82
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- CH <sub>3</sub> OH	1 : 2: 11mL	M(Ac) <sub>2</sub> ×2L	34,28/33,96	2,85/2,76	10,00/9,74
Ni(Ac) <sub>2</sub> ×4H <sub>2</sub> O-L- DMF	1 : 2: 11mL	M(Ac) <sub>2</sub> ×L×DMF	35,39/34,59	3,83/3,69	12,38/12,18

Fig. 1, a. Scheme of the proposed structure for the obtained complexes M(Ac)<sub>2</sub>·2LFig. 1, b. Scheme of the proposed structure for the obtained complexes M(Ac)<sub>2</sub>·L·DMF




Fig. 2. <sup>1</sup>H NMR of 5-methyl-3-(trifluoromethyl)-1H-pyrazole




Fig. 3. <sup>1</sup>H NMR of the mixture of the zinc complexes Zn(Ac)<sub>2</sub>·2H<sub>2</sub>O-L-Solv with 5-methyl-3-(trifluoromethyl)-1H-pyrazole




Fig. 4. IR spectra of the complexes  $M(\text{Ac})_2 \cdot 2L$  in the systems  $M(\text{Ac})_2 \cdot 4\text{H}_2\text{O} \cdot L \cdot \text{CH}_3\text{OH} / \text{CH}_3\text{CN}$  at a ratio of  $M:L = 1:1$




Fig. 5. IR spectra of the complexes  $M(\text{Ac})_2 \cdot 2L$  in the systems  $M(\text{Ac})_2 \cdot 4\text{H}_2\text{O} \cdot L \cdot \text{CH}_3\text{OH} / \text{CH}_3\text{CN}$  at a ratio of  $M:L = 1:2$




Fig. 6. IR spectra of the complexes  $M(\text{Ac})_2 \cdot L \cdot \text{DMF}$  in the systems  $M(\text{Ac})_2 \cdot 4\text{H}_2\text{O} \cdot L \cdot \text{DMF}$  at a ratio of  $M:L = 1:1$  and  $1:2$

*UV-Vis spectroscopy.* Diffuse reflection electron spectra were obtained for the synthesized compounds. Thus, the electronic spectrum of diffuse reflection for the copper powder sample of the obtained  $\text{Cu}(\text{Ac})_2 \cdot 2L$  complex (fig. 7)

showed the presence of a broad absorption band in the 550–700 nm region, which is attributed to the d-d transition and is characteristic of  $\text{Cu}^{2+}$  complexes.




Fig. 7. Electronic spectrum of diffuse reflection for the complex  $\text{Cu}(\text{Ac})_2 \cdot 2L$

The theoretical data for octahedral cobalt complexes are 400–500 nm (chromophore  $\text{CoO}_6\text{--CoN}_6$ ), while for tetrahedral complexes it is 600–700 nm (chromophore  $\text{CoN}_4$ ), the results obtained confirm the tetrahedral structure of the complex.

In the case of the cobalt (II) complex, a diffuse reflection electron spectrum was obtained, which showed a maximum

absorption in the region of  $12000 \text{ cm}^{-1}$ , indicating an octahedral coordination environment of the metal ion. Literature data for octahedral complexes indicate a transition from the  $\{\text{CoO}_6\}$  chromophore  $\sim 12000 \text{ cm}^{-1}$  to the  $\{\text{CoN}_6\}$  chromophore  $\sim 17000 \text{ cm}^{-1}$ , and a transition from the  $\{\text{CoO}_6\}$  chromophore ( $\sim 12455 \text{ cm}^{-1}$ ) to  $\{\text{CoN}_2\}$  ( $\sim 14770 \text{ cm}^{-1}$ ) (fig. 8).




Fig. 8. Electronic spectrum of diffuse reflection for the complex  $\text{Co}(\text{Ac})_2 \cdot 2\text{L}$

Therefore, it can be concluded that cobalt is in an octahedral nitrogen-oxygen donor environment.

The similarity in the structure of manganese and nickel complexes is evidenced by their diffuse reflection electron spectra. The presence of a broad band in the 450–550 nm

region, which is attributed to the d-d transition and is characteristic of manganese and nickel complexes. The bands in the 250 nm and 300 nm regions are attributed to charge transfer from the ligand to the metal (fig. 9).




Fig. 9. Electronic diffuse reflection spectra for  $\text{Mn}(\text{Ac})_2 \cdot 2\text{L}$  (top) and  $\text{Ni}(\text{Ac})_2 \cdot 2\text{L}$  (bottom) complexes

Therefore, in this paper, an organic ligand 5-methyl-3-(trifluoromethyl)-1H-pyrazole, was synthesized. The compound was identified by elemental analysis, IR and  $^1\text{H}$  NMR spectroscopy. The interaction of 3d metals such as Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) with 5-methyl-3-(trifluoromethyl)-1H-pyrazole was investigated. Methods for the synthesis of Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) compounds with 5-methyl-3-(trifluoromethyl)-1H-pyrazole were developed. Based on the data of elemental analysis and methods of IR, NMR, and electronic spectroscopy, the composition and structure of the obtained complexes have

been proposed. It has been shown that in the obtained complexes: the most typical method of coordination of the pyrazole cycle is realized: monodentate - through the pyridine nitrogen atom; the formation of two types of mononuclear Mn(II), Co(II), Ni(II), Cu(II), and Zn(II) complexes is observed:  $\text{M}(\text{Ac})_2 \cdot 2\text{L}$  and  $\text{M}(\text{Ac})_2 \cdot \text{L} \cdot \text{DMF}$ ; in the obtained complexes  $\text{M}(\text{Ac})_2 \cdot 2\text{L}$  and  $\text{M}(\text{Ac})_2 \cdot \text{L} \cdot \text{DMF}$  complexes, a six-coordinate environment of the central atom is realized due to four oxygen atoms from bidentate chelate coordinated acetate groups and two nitrogen atoms from molecules of a non-deprotonated ligand.

**Acknowledgements.** The author is grateful to the II European Chemistry School for Ukrainians for providing a comprehensive overview of current trends in European chemical science. <https://acmin.agh.edu.pl/en/detail/s/ii-european-chemistry-school-for-ukrainians>.

**Sources of funding.** This study did not receive any grant from a funding institution in the public, commercial, or non-commercial sectors. Funding provided by Taras Shevchenko National University of Kyiv.

#### References

- Annes, M. R. B., da Silva, M. A. P., & Soares, V. M. D. (2022). Recent advances on the use of fluorinated ligands in transition metal-catalyzed reactions. *Coordination Chemistry Reviews*, 455, 214368. <https://doi.org/10.1016/j.ccr.2021.214368>
- Davydenko, Y. (2025). Synthesis and Characterization of Coordination Compounds of Transition Metals Based on 5-Methyl-3-(Trifluoromethyl)-1H-Pyrazole. *ChemRxiv*. Preprint. <https://doi.org/10.26434/chemrxiv-2025-0brf5>
- Davydenko, Y. M., Vynohradov O. S., Gural'skiy I. A., Fritsky I. O., & Pavlenko V. O. (2024). *Coordination Compounds of Copper (II) with Pyrazole Ligands*. GRIN Verlag. <https://www.grin.com/document/1508817>
- Davydenko, Y. M., Vynohradov, O. S., Pavlenko, V. A., & Fritsky, I. O. (2023). Synthesis, structural and spectroscopic characterizations, hirshfeld surface analysis of two coordination compounds assembled from copper and carboxylates, 3, 5-dimethyl-1H-pyrazole. *Journal of Chemistry and Technologies*, 31(3), 468–476. <https://doi.org/10.15421/jchemtech.v31i3.276647>
- Davydenko, Y. M. (2011). Coordination compounds of 3d metals with non-chelating pyrazole ligands [Candidate dissertation of chemical sciences, Taras Shevchenko National University of Kyiv]. <https://ir.library.knu.ua/handle/123456789/6746>
- Davydenko, Y. M., Vynohradov, O. S., Pavlenko, V. O., Fesyeh, I. V., Fritsky, I. O., Naumova, D. D., & Shova, S. (2024). Synthesis, crystal structure, thermal stability and Hirshfeld surface analysis of azametallocrown copper(II) compounds with 4-iodopyrazole. *Journal of Chemistry and Technologies*, 32(3), 554–569. <https://doi.org/10.15421/jchemtech.v32i3.305413>
- Davydenko, Y. M., Vynohradov, O. S., Pavlenko, V. A., & Fritsky, I. O. (2025). Hirshfeld surface analysis of a mononuclear pyrazole-containing copper (II) complex obtained by oxidative dissolution method. *Journal of Chemistry and Technologies*, 33(2), 342–351. <https://doi.org/10.15421/jchemtech.v33i2.312363>
- Davydenko, Y. M., Vitske, V. A., Pavlenko, V. A., Haukka, M., Vynohradov, O. S., & Fritsky, I. O. (2022). Synthesis, crystal structure and properties of coordination polymers based on (3, 5-dimethyl-1H-pyrazole-4-yl)-acetic acid. *Journal of Chemistry and Technologies*, 30(2), 174–183. <https://doi.org/10.15421/jchemtech.v30i2.252517>
- Fustero, S., Simón-Fuentes, A., Delgado, O., & Román, R. (2014). Fluorinated Pyrazoles and Indazoles. In V. Nenajdenko (Ed.), *Fluorine in Heterocyclic Chemistry* (Vol. 1, pp. 279–321). Springer International Publishing. [https://doi.org/10.1007/978-3-319-04346-3\\_7](https://doi.org/10.1007/978-3-319-04346-3_7)
- Kieffer, M., Wally, T., Nguyen, T. K. T., Papper, V., Kathan, M., & Hecht, S. (2022). A unidirectional bipyrazolic photoswitch driven by an acido-catalyzed E/Z

- isomerization. *Journal of the American Chemical Society*, 144(25), 11470–11478. <https://doi.org/10.1021/jacs.2c04026>
- Kolarski, D., Feringa, B. L., Buma, W. J., & Szymanski, W. (2020). Visible-light-driven pyrazole photoswitches with near-quantitative photoconversion and high thermal stability. *Angewandte Chemie International Edition*, 59(32), 13398–13404. <https://doi.org/10.1002/anie.202003882>
- Li, Y., Wang, L., & Feng, S. (2023). Synthesis, crystal structure and magnetic properties of a new dinuclear copper(II) complex with 3-methyl-5-(trifluoromethyl)pyrazolate. *Inorganica Chimica Acta*, 555, 121585. <https://doi.org/10.1016/j.ica.2023.121585>
- Mezei, G., Zaleski, C. M., & Pecoraro, V. L. (2007). Structural and functional evolution of metallacrowns. *Chemical reviews*, 107(11), 4933–5003. <https://doi.org/10.1021/cr078200h>
- Mykhailiuk, P. K. (2020). Fluorinated pyrazoles: From synthesis to applications. *Chemical Reviews*, 121(3), 1670–1715. <https://doi.org/10.1021/acs.chemrev.0c01015>
- Pandolfo, L., & Pettinari, C. (2017). Trinuclear copper (ii) pyrazolate compounds: A long story of serendipitous discoveries and rational design. *CrystEngComm*, 19(13), 1701–1720.
- Park, H. S., Jeong, S. M., Han, J. T., & Lee, K. M. (2020). Fluorinated azopyrazoles as negative-type photoswitchable liquid crystal dopants. *ChemPhotoChem*, 4(6), 381–387. <https://doi.org/10.1002/cptc.202000030>
- Roy, S., Dutta, P. K., & Konar, S. (2023). A series of 3D metal-organic frameworks based on a pyrazole-dicarboxylate ligand: Synthesis, structure, and gas adsorption properties. *Inorganic Chemistry*, 62(1), 234–243. <https://doi.org/10.1021/acs.inorgchem.2c03381>
- Sánchez-Cantú, M. A., García-López, M., & Mendoza-Espinosa, D. (2018). The Role of Pyrazole-Based Ligands in Olefin Polymerization. *Coordination Chemistry Reviews*, 363, 1–24. <https://doi.org/10.1016/j.ccr.2018.02.012>
- Sun, L., Chen, Y., & Li, X. (2023). Luminescent lanthanide-organic frameworks constructed from a fluorinated pyrazole-carboxylate ligand for sensing of metal ions. *Crystal Growth & Design*, 23(8), 5678–5686. <https://doi.org/10.1021/acs.cgd.3c00424>
- Tití, A., Zaidi, K., Alzahrani, A. Y., El Kodadi, M., Yousfi, E. B., Moliterni, A., ... & Abboud, M. (2023). New in situ catalysts based on nitro functional pyrazole derivatives and copper (II) salts for promoting oxidation of catechol to o-quinone. *Catalysts*, 13(1), 162. <https://doi.org/10.3390/catal13010162>
- Vynohradov, O. S., Davydenko, Y. M., Pavlenko, V. A., Naumova, D. D., Shova, S., & Petlovanyi, D. (2022). Crystal structure and Hirshfeld surface analysis of dichloridotetrakis (4-methyl-1H-pyrazole-kN2) nickel (II) acetonitrile disolvate. *Structure Reports*, 78(12), 1156–1160. <https://doi.org/10.1107/S2056989022010362>
- Wang, X., Liu, J., & Gao, S. (2023). A family of tetranuclear manganese(II) pyrazolate complexes: Exploring magneto-structural correlations. *Chemistry – A European Journal*, 29(40), e202300654. <https://doi.org/10.1002/chem.202300654>
- Zhang, L., Wu, B., & Li, G. (2024). A luminescent 3D zinc(II) metal-organic framework based on a bis(pyrazolyl) ligand for selective sensing of Fe<sup>3+</sup> ions. *CrystEngComm*, 26(5), 724–731. <https://doi.org/10.1039/D3CE01234F>

Отримано редакцією журналу / Received: 10.09.25  
Прорецензовано / Revised: 01.10.25  
Схвалено до друку / Accepted: 13.10.25

Юлія ДАВИДЕНКО, канд. хім. наук

ORCID ID: 0000-0003-0223-0363

e-mail: y.davydenko@knu.ua

Київський національний університет імені Тараса Шевченка, Київ, Україна

## СИНТЕЗ І ХАРАКТЕРИСТИКА КООРДИНАЦІЙНИХ СПОЛУК ПЕРЕХІДНИХ МЕТАЛІВ НА ОСНОВІ 5-МЕТИЛ-3-(ТРИФТОРМЕТИЛ)-1Н-ПІРАЗОЛУ

**Вступ.** Синтез координаційних сполук на основі піразолів є надзвичайно важливим у сучасній хімії, оскільки ці сполуки мають універсальні властивості та різноманітне практичне застосування в таких сферах: каталіз, матеріалознавство, оптоелектроніка та медицина. Введення трифторметильних груп у такі ліганди є потужною стратегією для модуляції електронних і стеричних властивостей комплексів, що значно впливає на їхню кислотність, стабільність і реакційну здатність. У цьому контексті 5-метил-3-(трифторметил)-1H-піразол виділяється як особливо перспективний ліганд, координаційна хімія якого з перехідними металами залишається напролюд недослідженою. Незважаючи на потенціал 5-метил-3-(трифторметил)-1H-піразолу, систематичне дослідження синтезу, структурної різноманітності та властивостей його координаційних сполук в основному відсутнє в сучасній літературі. Ця прогалина є втраченою можливістю, оскільки унікальний електронний профіль цього ліганду може відкрити нові функціональні можливості в молекулярних матеріалах. Щоб розв'язати цю проблему, пропонується комплексне дослідження синтезу та характеристик нового сімейства координаційних сполук перехідних металів першого ряду (Mn, Co, Ni, Cu, Zn) з лігандом 5-метил-3-(трифторметил)-1H-піразол. Ця робота має на меті систематично дослідити його координаційний ландшафт, надаючи фундаментальні знання для майбутнього проєктування передових функціональних матеріалів.

**Методи.** Для ідентифікації отриманих сполук було використано різні методи, такі як: інфрачервона, електронна та ЯМР-спектроскопія. Усі матеріали були отримані з комерційних джерел без додаткового очищення. ІЧ-спектри синтезованих сполук записували на ІЧ-Фур'є-спектрометрі Perkin-Elmer BX (400–4000 см<sup>-1</sup>) у таблетках KBr. Електронні спектри дифузного відбиття отриманих зразків в УФ і видимому діапазоні реєстрували на спектрометрі Varian Cary 50. Інтенсивність вимірювали відносно KBr. Розчинники очищали за стандартними методиками. Спектри ЯМР <sup>1</sup>H записували на спектрометрі Bruker AC-400.

**Результати.** Взаємодія 5-метил-3-(трифторметил)-1H-піразолу (L) з ацетатами 3d-металів (Mn, Zn, Cu, Co, Ni) у неводних розчинах виявилася залежною від розчинника. У середовищі розчинників CH<sub>3</sub>CN і CH<sub>3</sub>OH (системи M(AC)<sub>2</sub>·4H<sub>2</sub>O–L–CH<sub>3</sub>OH/CH<sub>3</sub>CN) при співвідношенні M:L 1:1 і 1:2 утворювалися продукти зі складом M(AC)<sub>2</sub>·2L. Тим часом у диметилформаміді (системи M(AC)<sub>2</sub>·4H<sub>2</sub>O–L–DMF) за тих самих співвідношень були виділені комплекси M(AC)<sub>2</sub>·L·DMF, що містили молекулу розчинника. Склад сполук було підтверджено елементним аналізом, а також інфрачервоною, електронною та ЯМР-спектроскопією.

**В и с н о в к и .** Синтезовано органічний ліганд, 5-метил-3-(трифторметил)-1H-піразол. Сполука ідентифікувалась за допомогою елементного аналізу, ІЧ- та <sup>1</sup>H ЯМР-спектроскопії. Було досліджено взаємодію 3d-металів, таких як Mn(II), Co(II), Ni(II), Cu(II) і Zn(II), з 5-метил-3-(трифторметил)-1H-піразолом. Розроблено методи синтезу сполук Mn(II), Co(II), Ni(II), Cu(II) та Zn(II) з 5-метил-3-(трифторметил)-1H-піразолом. На основі даних елементного аналізу й методів ІЧ, ЯМР та електронної спектроскопії запропоновано склад і будову отриманих комплексів. Було показано, що в отриманих комплексах: реалізовано найбільш типовий спосіб координації піразольного циклу: монодентатний – через піридиновий атом нітрогену; отримано два типи моноядерних комплексів Mn(II), Co(II), Ni(II), Cu(II) та Zn(II): M(Ac)<sub>2</sub>·2L та M(Ac)<sub>2</sub>·L·DMF; в отриманих комплексах M(Ac)<sub>2</sub>·2L та M(Ac)<sub>2</sub>·L·DMF реалізується шестикоординаційне оточення центрального атома завдяки чотирьом атомам кисню від бідентатно-хелатно координованих ацетатних груп і двох атомів нітрогену від молекул недепротонованого ліганду.

**К л ю ч о в і с л о в а :** 5-метил-3-(трифторметил)-1H-піразол, комплекси, 3d-метали, <sup>1</sup>H ЯМР-спектроскопія, ІЧ-спектроскопія.

Автор заявляє про відсутність конфлікту інтересів. Спонсори не брали участі в розробленні дослідження; у зборі, аналізі чи інтерпретації даних, у написанні рукопису або у прийнятті рішення про публікацію результатів.

The author declares no conflicts of interest. The funders had no role in the design of the study, in the collection, analysis or interpretation of data, in the writing of the manuscript, or in the decision to publish the results.